



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **Alternating IV nerve palsy and ptosis as a first sign of childhood ocular myasthenia gravis**

Chappaz, A ; Knirsch, U ; Gerth-Kahlert, C

DOI: <https://doi.org/10.1016/j.pediatrneurol.2014.12.004>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-103703>

Journal Article

Accepted Version

Originally published at:


Chappaz, A; Knirsch, U; Gerth-Kahlert, C (2015). Alternating IV nerve palsy and ptosis as a first sign of childhood ocular myasthenia gravis. *Pediatric Neurology*, 52(4):460-461.

DOI: <https://doi.org/10.1016/j.pediatrneurol.2014.12.004>

# Accepted Manuscript

Alternating IV nerve palsy and ptosis as a first sign of childhood ocular myasthenia gravis

Ariane Chappaz, Ursula Knirsch, Christina Gerth-Kahlert



PII: S0887-8994(14)00761-9

DOI: [10.1016/j.pediatrneurol.2014.12.004](https://doi.org/10.1016/j.pediatrneurol.2014.12.004)

Reference: PNU 8562

To appear in: *Pediatric Neurology*

Received Date: 10 October 2014

Revised Date: 18 November 2014

Accepted Date: 23 December 2014

Please cite this article as: Chappaz A, Knirsch U, Gerth-Kahlert C, Alternating IV nerve palsy and ptosis as a first sign of childhood ocular myasthenia gravis, *Pediatric Neurology* (2015), doi: 10.1016/j.pediatrneurol.2014.12.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Alternating IV nerve palsy and ptosis as a first sign of childhood ocular myasthenia gravis**

Ariane Chappaz<sup>a</sup>, Ursula Knirsch<sup>b</sup> and Christina Gerth-Kahlert<sup>a</sup>

<sup>a</sup>University Hospital Zurich

Department of Ophthalmology

Frauenklinikstrasse 24

CH-8091 Zurich, Switzerland

<sup>b</sup>University Children's Hospital

Department of Pediatric Neurology

Steinwiesstrasse 75

CH-8032 Zurich, Switzerland

Corresponding author:

Christina Gerth-Kahlert

University Hospital Zurich, Switzerland

Department of Ophthalmology

Frauenklinikstrasse 24

CH-8091 Zurich

Phone +41-44-255 4940

FAX +41-44-255 2448

[christina.gerth-kahlert@usz.ch](mailto:christina.gerth-kahlert@usz.ch)

All authors have no conflict of interest.

The authors have not published or submitted any related papers about this case.

Word count: 500

A previously healthy five-year-old girl was referred due to diplopia of sudden onset. The patient presented with a moderate *right* ptosis, left inferior oblique overaction and superior oblique underaction, and a positive Bielschowsky head tilt test to the left. This suggested a *left* IV nerve palsy but the right ptosis remained undefined. (Fig.1A) Ophthalmic and neurological examination and cranial MRI were otherwise normal. 10 days later, a *right* IV nerve palsy with a *right* ptosis and left adduction deficit was evident. One month after initial presentation, a *left* ptosis and *left* IV nerve palsy were present. Simpson test was positive after 70sec and the ptosis improved after the ice pack test. Repetitive nerve stimulation of the facial nerve demonstrated an abnormal decrement. Serum acetylcholine receptor (AChR) antibody level was elevated. The diagnosis of ocular myasthenia gravis (OMG) was confirmed. A chest MRI excluded a thymoma. Treatment with pyridostigmine was initiated with a good response after an increase to 60 mg a day. Fourth nerve palsy resolved completely after six months and no generalization was observed.

OMG can mimic oculomotor disorders and should be considered in children with unusual and alternating oculomotor symptoms. Differential diagnoses are neoplasm, trauma, inflammatory or vascular causes. Ophthalmic signs such as ptosis or ophthalmoplegia present in about 90% of children with myasthenia with or without generalization.<sup>1</sup> In prepubertal children ocular symptoms appear mostly at disease onset and spontaneous remission is common. The risk for generalization decreases after two years of disease duration<sup>2</sup> but the development of generalized symptoms is still possible after a period of three years<sup>3</sup>. Untreated pediatric patients with OMG may have a relatively low risk of developing generalized symptoms<sup>2</sup>. Thymectomy, when performed before the onset of generalized symptoms, showed a trend toward protection from the development of generalized symptoms but no prospective data

are available.<sup>2</sup> AChR antibody level, although not always positive, repetitive nerve stimulation and, most sensitive (if the first remain negative) stimulated single fiber EMG is recommended to confirm the diagnosis. Pyridostigmine is the first line treatment of pediatric OMG. Treatment with steroids in young children is ambivalent due to the side effects and may not be necessary in pediatric OMG. Immunosuppressive agents may reduce the conversion of adult OMG to generalized myasthenia gravis but seems not to influence the course of the disease in children.<sup>2, 4</sup>

**Acknowledgement:**

We thank Prof. K. Landau, Dr. A. Klein and Mrs. R. Gulik for examination and discussion, Mrs. Stojcic and Mr. Breitschmid for the photographic work.

**References**

1. Andrews P. Autoimmune myasthenia gravis in childhood. *Semin Neurol* 2004; 24:101–110.
2. Pineles SL, Avery RA, Moss HE et al. Visual and Systemic Outcomes in Pediatric Ocular Myasthenia Gravis. *Am J Ophthalmol*. 2010; 150:453–459.
3. Marina AD, Trippe H, Lutz S, Schara U. Juvenile Myasthenia Gravis: Recommendations for Diagnostic Approaches and Treatment. *Neuropediatrics* 2014; 45:75–83.
4. Ortiz S, Borchert M. Long-term Outcomes of Pediatric Ocular Myasthenia Gravis. *Ophthalmology* 2008; 115:1245–1248.

**Figure:**

Presentations with alternating sides of ptosis and 4<sup>th</sup> nerve palsy.



